

Analytic proof of the Weinberg theorem for off mass shell and composite pions

Poster for the 5th "Quark confinement and the hadron spectrum" Conference, Gargnano, September 2002
Pedro Bicudo

Departamento de Física, and CFIF, Instituto Superior Técnico, Av. Rovisco Pais, 1049-001 Lisboa, Portugal

The Weinberg theorem for the pion-pion scattering is proved analytically in full detail for quark models in the ladder truncation. The proof is displayed with Feynman diagrams. The axial and vector Ward identities, for the quark propagator and for the ladder, exactly cancel any model dependence. Off mass shell and finite size effects are included in the quark-antiquark pion Bethe Salpeter vertices. This is applied to compute the isospin 0, 1 and 2 scattering matrices off the mass shell.


The success of the Quark Model [1] relies on its ability to reproduce the Hadron spectrum with microscopical interacting quarks. This includes for instance the linear Regge trajectories, and the correct spin-spin and spin-orbit interactions. Moreover the quark model is competent to predict microscopically the hadron-hadron interactions [2]. For recent coupled channel studies see [3,4,5]. However the quark model suffered from initial difficulties of fully understanding the low pion mass and the small π - π scattering length. With the aim to cure this important problem, chiral symmetry breaking was introduced in the quark model. [6,7,8]. This opened the possibility to extend the low energy pion theorems of current algebra to the quark model framework.

In this paper we prove that the $\pi - \pi$ scattering theorem of Weinberg [9] applies to quark models with chiral invariant quark-quark interactions, including the Nambu and Jona-Lasinio model [10,11], conventional quark models with confining instantaneous interactions [4,5], models with Euclidean space integrations [12,5,13], and also covariant models in Minkowsky space [14]. After the original work of Weinberg [9], the theorem was also derived with Ward identities for the pion fields [15] and with a functional integration of quarks [16]. However these methods use effective pions, whereas pions in quark models have naturally a finite size [17] which might affect $\pi - \pi$ scattering.

Here we complete in full detail an analytical proof which was recently outlined in [5], using Ward identities and the diagrammatic method [18]. Moreover we extend the proof to pions with off mass shell momenta. Amplitudes for off mass shell pions are crucial because $\pi - \pi$ scattering is experimentally estimated from $\pi - N$ scattering. The nucleon N provides a virtual off mass shell pion π^* that scatters with the incident π . For instance a virtual pion π^* with offshellness $P^2 - M_\pi^2 = -3.32M_\pi^2$ is present in the π production at threshold.

To prove the Weinberg theorem in the Quark Model formalism we have to truncate the series of Feynman di-

agrams consistently. The ladder truncation is commonly assumed for the boundstate equations,




$$\text{Box} = \text{Line} + \text{Box with internal lines} \quad (1)$$

where the dotted line corresponds to the chiral invariant quark-quark interaction of vertex V and kernel $\mathcal{K}$. The arrowed line corresponds to the Dirac quark propagator. The Ward identities [19] show that the ladder is consistent with the quark self energy equation in the rainbow approximation. When chiral symmetry is spontaneously broken, the dressed quark propagator is non trivial,

$$S = \frac{i}{A(k^2) \not{k} - B(k^2)}, \quad (2)$$

and the π is a Goldstone boson in the chiral limit. There is also evidence that the hadron-hadron coupled channel equations should include one meson exchange, both in the sigma model [9,20], in the Nambu and Jona-Lasinio model [11], in the constituent quark models [5], and in an Euclidean quark model [5]. In microscopic calculations the Feynman diagram for $\pi - \pi$ scattering must therefore include inside the box a vertical scalar ladder and a horizontal scalar ladder, see paper [5] for details. To cure double counting we also have to subtract an empty box.

The main task of this paper consists in computing *independently of the Quark Model*, and up to order M_π^2 and $P_i P_j$ in the π mass and momenta, the Feynman loop,




$$\text{Box} + \text{Box} - \text{Box} \quad (3)$$

where χ is the Bethe Salpeter vertex of the pion. The subindex P_i accounts an external momentum flowing into the loop.

To derive the proof it is convenient to define the dressed axial vertex Γ_A with the axial Ward identity,

$$\Gamma_A(k_1, k_2) = S^{-1}(k_1)\gamma_5 + \gamma_5 S^{-1}(k_2), \quad (4)$$

because it is related to the pion vertex and because it has the crucial property,



$$\text{Box} = \text{Box} + \text{Box} \quad (5)$$

which constitutes a Ward identity for the ladder. This identity is derived if we expand the ladders and substitute

the vertex in the left hand side. Then all terms with an intermediate γ_5 include the anticommutator $\{\gamma_5, V\}$ and this cancels because *the interaction is chiral invariant* and *the kernel is local*. Only the right hand side survives.

We now derive a second useful relation for Γ_A . The Dyson Schwinger equation for the full propagator is,

$$S(k)^{-1} = S_0^{-1}(k) - \int \frac{d^4 q}{(2\pi)^4} \mathcal{K}(q) V S(k+q) V \quad (6)$$

where V is some vertex that anticommutes with γ_5 and that is color dependent as well, $\mathcal{K}$ is the quark-quark Kernel, and $S_0 = i/(\not{k} + m)$ is the bare quark propagator. If we multiply right or left with γ_5 and sum, we get,

$$S(k_1)^{-1} \gamma_5 + \gamma_5 S(k_2)^{-1} = S_0(k_1)^{-1} \gamma_5 + \gamma_5 S_0(k_2)^{-1} + \int \mathcal{K}(q) V (S(k_1 + q) \gamma_5 + \gamma_5 S(k_2 + q)) V, \quad (7)$$

and this is the Bethe Salpeter equation for the vertex,

$$\begin{aligned} \Gamma_A(k_1, k_2) &= \gamma_A(k_1, k_2) + \int \mathcal{K}(q) V S(k_1 + q) \Gamma_A S(k_2 + q) V \\ \gamma_A(k_1, k_2) &= S_0^{-1}(k_1) \gamma_5 + \gamma_5 S_0^{-1}(k_2), \end{aligned} \quad (8)$$

where the particular part γ_A is the bare axial vertex,

$$\gamma_A = \frac{(\not{P} - 2m)}{i} \gamma_5, \quad P = k_1 - k_2. \quad (9)$$

γ_A vanishes when the current quark mass m is small (chiral limit) and at the same time the total momentum P^μ of the vertex is small. On the other hand the dressed vertex Γ_A is finite,

$$\Gamma_A(k_1, k_2) = \frac{A(k_1) \not{k}_1 - A(k_2) \not{k}_2 - B(k_1) - B(k_2)}{i} \gamma_5, \quad (10)$$

providing spontaneous chiral symmetry breaking occurs in eq. (6) to generate a dynamical mass in the dressed quark propagator. After we iterate the Bethe Salpeter equation for the dressed axial vertex Γ_A , and include the external propagators, we get the second useful relation,

$$S \Gamma_A S = \text{diagram} \rightarrow \gamma_A. \quad (11)$$

The main step to get the proof consists in decreasing the number of vertices with the help of eqs. (5) and (11),

$$\begin{aligned} \text{diagram} &= \text{diagram} \rightarrow \text{diagram} \\ &= \gamma_5 \text{diagram} + \text{diagram} \rightarrow \gamma_5 \gamma_A. \end{aligned} \quad (12)$$

This is repeated to compute the square box with ladder,

$$\begin{aligned} \text{diagram} &= \frac{1}{2} \text{diagram} \\ &+ 2 \{ \gamma_A, \gamma_5 \} \text{diagram} + \frac{1}{4} \{ \gamma_A, \gamma_5 \} \text{diagram} \{ \gamma_A, \gamma_5 \}, \end{aligned} \quad (13)$$

and we get three terms, respectively of order 1, γ_A and γ_A^2 . We note that all the other factors (Γ_A , S , and the scalar and vector ladder) are finite and carry the scale of the effective quark-quark interaction.

When we sum the three diagrams of the full box,

$$\begin{aligned} \text{diagram} &+ \text{diagram} - \text{diagram} \\ &= 8 i m \text{tr}\{S\} + o(m^2, P_i^4, m P_i^2), \end{aligned} \quad (14)$$

the finite term of order 1 *exactly cancels*. The total box vanishes in the chiral limit of $m = 0$, although each one of the three boxes is generally finite and proportional to the fourth power of the scale of the interaction. This cancellation in eq. (15) complies with the Adler zero and corresponds to a higher Ward identity.

The dependence of eq. (15) on the quark current mass m is trivially computed with eqs. (9) and (13). The exact cancellation in the momentum dependence is more subtle. Expanding up the first order in $P_i P_j$, the γ_A and γ_A^2 terms in eq. (13),

$$\frac{P_3^\mu - P_4^\mu}{i} \left[-i \partial_\mu S^{-1} \text{diagram} + \gamma_\mu \text{diagram} \right] \rightarrow \frac{P_1 - P_2}{i} \quad (16)$$

and this cancels due to the vector Ward Identity which has and analogous variant to eq. (11).

We now apply these Ward identity techniques to the actual Feynman amplitude with pion vertices of eq. (3). In the limit of vanishing current quark mass m and vertex momentum P^μ , the Bethe-Salpeter (8) equation for the axial vertex Γ_A becomes homegenous and is thus identical to the homogeneous Bethe-Salpeter equation for the pion vertex $\chi_{\pi P}(k)$. In this limit the pion is a massless Goldstone boson, and the pion Bethe Salpeter vertex is proportional to the dressed axial vertex, and to the dynamical quark mass $B(k)$,

$$\chi_0(k) = \frac{B(k)}{n_\pi} \gamma_5 = \frac{1}{2 i n_\pi} \Gamma_{A0}(k, k), \quad (17)$$

where n_π is the norm. And the loop (3) *vanishes*.

To get the loop (3) up to order P_i^2 and M_π^2 , we need at most two full Bethe Salpeter vertices χ , the other two can be approximated by $\Gamma_A/(2 i n_\pi)$. Expanding up to second order in $\chi - \Gamma_A/(2 i n_\pi)$ and regrouping the sum, we find that the amplitude of eq.(3) is the sum of four classes of terms. Each class includes a sum of the possible cyclic permutations of the external momenta P_1, P_2, P_3 , and P_4 . We get, eq.(14) with factor 3 times $(2 i n_\pi)^{-4}$, minus, with factor 2 times $(2 i n_\pi)^{-3}$, (4 permutations)

$$\text{diagram} + \text{diagram} - \text{diagram} \quad (18)$$

plus, with factor $(2i n_\pi)^{-2}$, (4 permutations)

$$\begin{array}{c} \chi_{P_4} \uparrow \downarrow \chi_{P_1} \\ \text{[Diagram: A box with four external lines. Top-left: incoming \chi_{P_4}, top-right: incoming \chi_{P_1}, bottom-left: outgoing \Gamma_{AP_3}, bottom-right: outgoing \Gamma_{AP_2}.]}\end{array} + \begin{array}{c} \chi_{P_4} \uparrow \downarrow \chi_{P_1} \\ \text{[Diagram: A box with four external lines. Top-left: incoming \chi_{P_4}, top-right: incoming \chi_{P_1}, bottom-left: outgoing \Gamma_{AP_3}, bottom-right: outgoing \Gamma_{AP_2}.]}\end{array} - \begin{array}{c} \chi_{P_4} \uparrow \downarrow \chi_{P_1} \\ \text{[Diagram: A box with four external lines. Top-left: incoming \chi_{P_4}, top-right: incoming \chi_{P_1}, bottom-left: outgoing \Gamma_{AP_3}, bottom-right: outgoing \Gamma_{AP_2}.]}\end{array}, \quad (19)$$

plus, with factor $(2i n_\pi)^{-2}$, (2 permutations)

$$\begin{array}{c} \chi_{P_4} \uparrow \downarrow \Gamma_{AP_1} \\ \text{[Diagram: A box with four external lines. Top-left: incoming \chi_{P_4}, top-right: incoming \Gamma_{AP_1}, bottom-left: outgoing \Gamma_{AP_3}, bottom-right: outgoing \chi_{P_2}.]}\end{array} + \begin{array}{c} \chi_{P_4} \uparrow \downarrow \Gamma_{AP_1} \\ \text{[Diagram: A box with four external lines. Top-left: incoming \chi_{P_4}, top-right: incoming \Gamma_{AP_1}, bottom-left: outgoing \Gamma_{AP_3}, bottom-right: outgoing \chi_{P_2}.]}\end{array} - \begin{array}{c} \chi_{P_4} \uparrow \downarrow \Gamma_{AP_1} \\ \text{[Diagram: A box with four external lines. Top-left: incoming \chi_{P_4}, top-right: incoming \Gamma_{AP_1}, bottom-left: outgoing \Gamma_{AP_3}, bottom-right: outgoing \chi_{P_2}.]}\end{array}. \quad (20)$$

To repeat the main step of eq. (12), replacing one of the two Γ_{AP} by a χ_P , it is convenient to include (or remove) a ladder in the pion vertex χ_P . In the neighborhood of a boundstate b pole M_b^2 in the external momentum P^2 ,

$$\begin{array}{c} \text{[Diagram: A box with four external lines. Top-left: incoming \chi_{P_4}, top-right: incoming \chi_{P_1}, bottom-left: outgoing \Gamma_{AP_3}, bottom-right: outgoing \Gamma_{AP_2}.]}\end{array} = \begin{array}{c} \text{[Diagram: A box with four external lines. Top-left: incoming \chi_{P_4}, top-right: incoming \chi_{P_1}, bottom-left: outgoing \Gamma_{AP_3}, bottom-right: outgoing \Gamma_{AP_2}.]}\end{array} \chi_{b_P} \frac{i}{P^2 - M_b^2 + i\epsilon} \chi_{b_{-P}} \begin{array}{c} \text{[Diagram: A box with four external lines. Top-left: incoming \chi_{P_4}, top-right: incoming \chi_{P_1}, bottom-left: outgoing \Gamma_{AP_3}, bottom-right: outgoing \Gamma_{AP_2}.]}\end{array} \quad (21)$$

multiplying the right of both sides with the vertex χ ,

$$\begin{array}{c} \text{[Diagram: A box with four external lines. Top-left: incoming \chi_{P_4}, top-right: incoming \chi_{P_1}, bottom-left: outgoing \Gamma_{AP_3}, bottom-right: outgoing \Gamma_{AP_2}.]}\end{array} \chi = \begin{array}{c} \text{[Diagram: A box with four external lines. Top-left: incoming \chi_{P_4}, top-right: incoming \chi_{P_1}, bottom-left: outgoing \Gamma_{AP_3}, bottom-right: outgoing \Gamma_{AP_2}.]}\end{array} \chi \frac{P^2 - M_\pi^2}{i\mathcal{I}}, \quad (22)$$

$$\mathcal{I} = \chi \begin{array}{c} \text{[Diagram: A box with four external lines. Top-left: incoming \chi_{P_4}, top-right: incoming \chi_{P_1}, bottom-left: outgoing \Gamma_{AP_3}, bottom-right: outgoing \Gamma_{AP_2}.]}\end{array} \chi, \quad (22)$$

where $\mathcal{I}$ is finite and proportional to the square of the scale of the interaction, nevertheless it will factorize from the results. Moreover the off mass shell Bethe Salpeter equation is derived replacing eq. (22) in eq. (1),

$$\chi_{P=} \begin{array}{c} \text{[Diagram: A box with four external lines. Top-left: incoming \chi_{P_4}, top-right: incoming \chi_{P_1}, bottom-left: outgoing \Gamma_{AP_3}, bottom-right: outgoing \Gamma_{AP_2}.]}\end{array} \chi_P \left(1 - \frac{P^2 - M_\pi^2}{i\mathcal{I}}\right)^{-1}. \quad (23)$$

The off mass shell correction only contributes to the Bethe Salpeter equation (23) at the order P^2 and M^2 . Therefore, up to the first order in the external P^μ , the vertex χ_P is *formally the same function* of P^μ , both for mass shell and for off mass shell pions.

The diagrams eventually simplify into traces related to the pion decay relation [21],

$$\text{tr}\{(S\chi S)_P \gamma^\mu \gamma^5\} = 2f_\pi P^\mu, \quad (24)$$

which measures f_π in the electroweak decay of pions on the mass shell. Because off mass shell vertices χ_P are formally the same as the mass shell ones, eq. (24) is also correct for $P^2 \neq M^2$. This is experimentally confirmed with the Goldberger Treiman relation [22]. Another useful trace is directly extracted, substituting the spectral decomposition of the ladder (21) in eq. (11),

$$\text{tr}\{(S\chi S)_{-P} \gamma_{AP}\} = 2n_\pi (P^2 - M_\pi^2). \quad (25)$$

A detailed momentum analysis of eqs (24,25) shows that the norm $n_\pi = i f_\pi$ and produces the desired traces,

$$\text{tr}\{(S\chi S)_{P_1} \gamma_{AP_2}\} = -2n_\pi (P_1 \cdot P_2 + M_\pi^2), \quad (26)$$

$$-m\text{tr}\{S\} = n_\pi^2 M_\pi^2, \quad (27)$$

including the Gell-Mann Oakes and Renner relation [23].

This completes the necessary steps to compute the loop (3). The diagrams of eq. (14) are trivial to compute, completing eq. (15) with the Gell-Mann Oakes and Renner relation (27). The three other cases (18,19,20) are now straightforwardly computed with the techniques that lead to eq. (13), replacing eq. (11) by eq. (22). It is clear that when we replace a γ_A by $\chi(P_j^2 - M_\pi^2)/i\mathcal{I}$, the order in P_j increases. Therefore the trace that replaces the term with a pair of $\{\gamma_A, \gamma_5\}$ in eq. (13) now vanishes up to the order of $P_i P_j$ and of M_π^2 . The trace that replaces the term of order $\{\gamma_A, \gamma_5\}$ in eq. (13) is now computed with the trace (26). Summing the total contributions to the Feynman loop of eq. (3) we get,

$$\begin{aligned} & + 3 \left(\frac{1}{2i n_\pi}\right)^4 (-8i n_\pi^2 M_\pi^2) \\ & - 2 \left(\frac{1}{2i n_\pi}\right)^3 \sum_{4\text{ perm.}} 2n_\pi (P_1^2 + P_1 \cdot P_3 - 2M_\pi^2) \\ & + \left(\frac{1}{2i n_\pi}\right)^2 \sum_{4\text{ perm.}} \frac{P_1^2 + P_2^2 + P_1 \cdot P_2 + P_1 \cdot P_3 + P_2 \cdot P_4 - 2M_\pi^2}{i} \\ & + \left(\frac{1}{2i n_\pi}\right)^2 \sum_{2\text{ perm.}} \frac{P_1^2 + P_3^2 - 2M_\pi^2}{i} \\ & = \frac{i}{4f_\pi^2} [(P_1 + P_2)^2 + (P_1 + P_4)^2 - 2M_\pi^2], \quad (28) \end{aligned}$$

where the conservation $P_1 + P_2 + P_3 + P_4 = 0$ of momentum was used to simplify the result.

We finally compute the $\pi - \pi$ scattering matrix T . We simply have to match the external pions, $i1$ and $i2$ incoming and $o1$ and $o2$ outgoing, with the four pion vertex that we just computed in eq.(28),

$$\begin{array}{ccc} \leftarrow q_{o1}, \tau_{o1} & P_4 \searrow \swarrow P_1 & q_{i1}, \tau_{i1} \leftarrow \\ \text{OUT} & \bullet & \text{IN} \\ \leftarrow q_{o2}, \tau_{o2} & P_3 \nearrow \nwarrow P_2 & q_{i2}, \tau_{i2} \leftarrow \end{array} \quad (29)$$

where the loop (3) is represented by the full circle. The loop is topologically invariant for cyclic permutations of P_1, P_2, P_3 and P_4 . To remove double counting we fix one match, say $P_1 = q_{i1}$. Then there are six different combinations of the remaining external legs.

In what concerns color, all the combinations are identical because the pion is a color singlet, and the color factor is appropriately included in the definition of f_π , see eq. (24). In what concerns momentum, we express our result in the usual Mandelstam relativistic invariant variables s, t and u . For instance the direct combination in eq. (29) produces the result $i(s + u - 2M_\pi^2)/(4f_\pi^2)$. We now introduce flavor. For simplicity it was not regarded in the previous definitions of the vertices Γ_{AP} and χ_P . Because the pion is an isovector, we have three different cases $I = 0, I = 1$ and $I = 2$. The flavor contributions

to the pion vertex simply factorize from the momentum contribution, and the different combinations only produce two classes of flavor traces, see Table I. Summing the six possible combinations of color, spin, momentum and flavor traces, and dividing by $-i$, we finally get the $\pi - \pi$ scattering T^I matrix,

$$\begin{aligned} T^0 &= -\frac{2s-M_\pi^2}{2f_\pi^2} - \frac{s+t+u-4M_\pi^2}{2f_\pi^2}, \\ T^1 &= -\frac{t-u}{2f_\pi^2}, \\ T^2 &= -\frac{-s+2M_\pi^2}{2f_\pi^2} - \frac{s+t+u-4M_\pi^2}{2f_\pi^2}, \end{aligned} \quad (30)$$

where $s+t+u-4M_\pi^2$ expresses the off mass shell contribution. We remark that eq. (30) complies with the Gasser and Leutwyler results. [15]. The T^I matrices of eq. (30) are computed at the tree level (including scalar and vector s, t, and u exchange), which is exact up to the order of $P_i^2 P_j^2$ and of M_π^2 .

The $\pi - \pi$ mass shell scattering lengths a_0^I are simply obtained from the scattering amplitudes T^0, T^2 with the Born factor of $\frac{-1}{16\pi M_\pi}$, and for vanishing 3-momenta. The $I = 1$ case is antisymmetric so the first scattering parameter is a_1^1 and the corresponding factor is $\frac{-1}{16\pi M_\pi} \frac{4}{3(t-u)}$,

$$\begin{aligned} a_0^0 &= \frac{7}{32\pi} \frac{M_\pi}{f_\pi^2}, \\ a_1^1 &= \frac{1}{24\pi} \frac{1}{M_\pi f_\pi^2}, \\ a_0^2 &= \frac{-1}{16\pi} \frac{M_\pi}{f_\pi^2}, \end{aligned} \quad (31)$$

this is the result of the famous Weinberg theorem for $\pi - \pi$ scattering [9]. Off mass shell effects are very important for the experiments. For instance in the scattering at $\pi - \pi$ threshold of a π beam with virtual π^* provided by a nucleon target, we estimate that the off mass shell effects of eq. (30) decrease a_0^0 by a factor of 0.5 and increase a_0^2 by a factor of 1.7.

We conclude that the Weinberg theorem for pion-pion low energy scattering is also exact when the pions are off the mass shell and have a finite size. It is remarkable that the current quark mass m and the traces $tr\{S\}$, $tr\{(S\Gamma_A)^4\}$, $tr\{(S\chi)^2\}$, which depend on the finite scale of the interaction, say the string tension σ or Λ_{QCD} , disappear from the final result. Any quark model with a chirally symmetric interaction complies with the Weinberg theorem. This detailed proof confirms that Goldstone bosons non only are massless but also are non-interacting at low energy.

I acknowledge Emilio Ribeiro for reporting on non-trivial Ward identities. I also acknowledge that discussions with the collaborators of the recent works [3,4,5] motivated the search for an analytical proof of the Weinberg theorem and identified the class of diagrams that should be included in the box of eq. (3).

-
- [1] A. De Rujula, H. Georgi and S. L. Glashow, Phys. Rev. D **12**, 147 (1975).
 - [2] J. Ribeiro, Z. Phys. C **5**, 27 (1980).
 - [3] P. Bicudo, G. Krein and J. Ribeiro, Phys. Rev. C **64**, 025202 (2001) [arXiv:hep-ph/0105289];
 - [4] P. Bicudo, M. Faria, G. Marques, and J. Ribeiro [arXiv:nucl-th/0106071].
 - [5] P. Bicudo, S. Cotanch, F. Llanes-Estrada, P. Maris, E. Ribeiro and A. Szczepaniak, Phys. Rev. D **65**, 076008 (2002) [arXiv:hep-ph/0112015].
 - [6] A. Le Yaouanc, L. Oliver, O. Pene and J.-C. Raynal, Phys. Rev. D **29**, 1233 (1984); Phys. Rev. D **31**, 137 (1985).
 - [7] S. Adler and A. Davis, Nucl. Phys. B **244**, 469 (1984).
 - [8] P. Bicudo and J. Ribeiro, Phys. Rev. D **42**, 1611 (1990); Phys. Rev. D **42**, 1625 (1990); Phys. Rev. D **42**, 1635 (1990).
 - [9] S. Weinberg, Phys. Rev. Lett. **17**, 616 (1966).
 - [10] Y. Nambu and J. Jona-Lasinio, Phys. Rev. **124**, 246 (1961); Phys. Rev. **122**, 345 (1961).
 - [11] V. Bernard, U. G. Meissner, A. Blin and B. Hiller, Phys. Lett. **B253**, 443 (1991); V. Bernard, A. H. Blin, B. Hiller, Y. P. Ivanov, A. A. Osipov and U. Meissner, Annals Phys. **249**, 499 (1996) [arXiv:hep-ph/9506309].
 - [12] Y. Dai, C. Huang and D. Liu, Phys. Rev. D **43**, 1717 (1991).
 - [13] S. Cotanch and P. Maris, [arXiv:hep-ph/0210151].
 - [14] V. Sauli, [arXiv:hep-ph/0108160].
 - [15] J. Gasser and H. Leutwyler, Annals Phys. **158**, 142 (1984).
 - [16] C. Roberts, R. Cahill, M. Sevier and N. Iannella, Phys. Rev. D **49**, 125 (1994) [arXiv:hep-ph/9304315].
 - [17] S. R. Amendolia *et al.*, Phys. Lett. B **146**, 116 (1984), E. B. Dally *et al.*, Phys. Rev. Lett. **48**, 375 (1982).
 - [18] P. Bicudo, Phys. Rev. C **60**, 035209 (1999) [arXiv:nucl-th/9802058].
 - [19] P. Bicudo, D.-S. Liu, J. Ribeiro, J. Villate Phys. Rev. D **47**, 1145 (1993).
 - [20] V. De Alfaro, S. Fubini, G. Furlan, C. Rossetti, "Currents in Hadron Physics", Amsterdam, North-Holland, (1973).
 - [21] W. Marciano and H. Pagels, Phys. Rept. **36**, 137 (1978).
 - [22] M. L. Goldberger and S. B. Treiman, Phys. Rev. **111**, 354 (1958).
 - [23] M. Gell-Mann, R. J. Oakes and B. Renner, Phys. Rev. **175**, 2195 (1968).

TABLE I. Table of the flavor traces. $tr\{\tau_{i1}\tau_{i2}\tau_{o2}^\dagger\tau_{o1}^\dagger\}$ and $tr\{\tau_{i1}\tau_{o2}^\dagger\tau_{i2}\tau_{o1}^\dagger\}$ are examples of the direct and exchange cases.

$I m_I$	$\tau_{i1}\tau_{i2}$	direct	exchange
00	$\frac{\vec{\sigma}\cdot\vec{\sigma}}{2\sqrt{3}}$	$\frac{3}{2}$	$-\frac{1}{2}$
11	$\frac{\sigma_1\sigma_2-\sigma_2\sigma_1}{2\sqrt{2}}$	1	0
22	$\sigma^+\sigma^+$	0	1